

HYPERVELOCITY STARS II. THE BOUND POPULATION

WARREN R. BROWN, MARGARET J. GELLER, SCOTT J. KENYON, MICHAEL J. KURTZ
Smithsonian Astrophysical Observatory, 60 Garden St, Cambridge, MA 02138

AND

BENJAMIN C. BROMLEY

Department of Physics, University of Utah, 115 S 1400 E, Rm 201, Salt Lake City, UT 84112

Accepted to ApJ

ABSTRACT

Hypervelocity stars (HVSs) are stars ejected completely out of the Milky Way by three-body interactions with the massive black hole in the Galactic center. We describe 643 new spectroscopic observations from our targeted survey for HVSs. We find a significant (3.5σ) excess of B-type stars with large velocities $+275 < v_{rf} < 450$ km s⁻¹ and distances $d > 10$ kpc that are most plausibly explained as a new class of HVSs: stars ejected from the Galactic center on bound orbits. If a Galactic center ejection origin is correct, the distribution of HVSs on the sky should be anisotropic for a survey complete to a fixed limiting apparent magnitude. The unbound HVSs in our survey have a marginally anisotropic distribution on the sky, consistent with the Galactic center ejection picture.

Subject headings: Galaxy: halo — Galaxy: center — Galaxy: stellar content — Galaxy: kinematics and dynamics — stars: early-type

1. INTRODUCTION

HVSs are a natural consequence of a massive black hole (MBH) in a dense stellar environment like that in the Galactic center. Hills (1988) first pointed out that a stellar binary encountering the Milky Way's central MBH can eject one member of the binary as a HVS traveling at $\sim 1,000$ km s⁻¹. HVSs differ from classical “runaway” stars because 1) HVSs are unbound and 2) the classical supernova ejection (Blaauw 1961) and dynamical ejection (Poveda et al. 1967) mechanisms that explain runaway stars cannot produce ejection velocities larger than 200 - 300 km s⁻¹ for B-type stars (Leonard 1991, 1993; Portegies Zwart 2000; Davies et al. 2002; Gualandris et al. 2004; Dray et al. 2005). The first HVS discovered, by comparison, has a heliocentric radial velocity of $+853$ km s⁻¹ and a Galactic rest-frame velocity of at least $+709 \pm 12$ km s⁻¹ (Brown et al. 2005a), many times that needed to escape the Milky Way. Photometric follow-up shows that the object is a slowly pulsating B main sequence star (Fuentes et al. 2006). Only interaction with a MBH can plausibly accelerate this 3 M_⊙ main sequence B star to such an extreme velocity and distance (~ 110 kpc).

HVSs are fascinating because they can be used to understand the nature and environs of MBHs (see the recent theoretical work of Gualandris et al. 2005; Holley-Bockelmann et al. 2005; Ginsburg & Loeb 2006a; Levin 2006; Perets et al. 2006; Baumgardt et al. 2006; Sesana et al. 2006; Bromley et al. 2006; Merritt 2006; Demarque & Virani 2007; Ginsburg & Loeb 2006b; O’Leary & Loeb 2006; Gualandris & Portegies Zwart 2007; Kollmeier & Gould 2007). The trajectories of HVSs also provide unique probes of the shape and orientation of the Galaxy’s dark matter halo (Gnedin et al. 2005). Discoveries of additional HVSs (Edelmann et al. 2005; Hirsch et al. 2005; Brown et al. 2006a,b) are starting to provide suggestive limits on the stellar mass function of HVSs,

the origin of massive stars in the Galactic Center, and the history of stellar interactions with the MBH. Clearly, a larger sample of HVSs will be a rich source for further progress on these issues.

We have designed a successful targeted survey for new HVSs. We use the 6.5m MMT and the Whipple 1.5m Tillinghast telescopes to obtain radial velocities of faint B-type stars, stars with lifetimes consistent with travel times from the Galactic center but which should not otherwise exist in the distant halo. Four earlier HVS discoveries from this survey are published elsewhere (Brown et al. 2006a,b). Here, we present spectroscopic observations of 643 new HVS candidates.

Our paper is organized as follows. In §2 we discuss our target selection and spectroscopic identifications. In §3 we present evidence for stars ejected from the Galactic center on bound orbits. In §4 we show that the unbound HVSs have a marginally anisotropic distribution on the sky. We conclude in §5. Our new observations are listed in Appendix A.

2. DATA

2.1. Target Selection

Brown et al. (2006b) describe our target selection for candidate HVSs. Briefly, we use Sloan Digital Sky Survey (SDSS) photometry to select late B-type stars by color. Figure 1 presents the HVS survey color-color selection. We use the 6.5m MMT telescope to observe every star with $17 < g'_0 < 19.5$ in this color range. We have observed 136 new HVS candidates and have completed the high declination region of the SDSS Data Release 4 (DR4, Adelman-McCarthy et al. 2006). The MMT survey is now 80% complete across SDSS DR4 and covers an area of $\sim 5,000$ deg² or 12% of the entire sky.

We have also carried out a new, complementary HVS survey targeting bright stars with $15 < g'_0 < 17$ and B-type colors in the SDSS. We use the 1.5m Tillinghast telescope and the FAST spectrograph (Fabricant et al. 1998) to observe these bright HVS candidates.




FIG. 1.— Color-color diagram showing the target selection regions for the MMT (*dashed line*) and FAST (*solid line*) samples, and the spectroscopic identifications for the combined set of 894 HVS candidates. The brighter FAST sample ($15 < g'_0 < 17$) targets bluer stars and finds a number of stars with B3 and B4 spectral types (*solid dots*).

We are able to survey efficiently much bluer stars with FAST because contamination from white dwarfs is less problematic at bright magnitudes. Figure 1 shows the FAST color-color selection: $0.1 < (u' - g')_0 < 1.5$ and $(10.67(g' - r')_0 + 3.5) < (u' - g')_0 < (10.67(g' - r')_0 + 5.07)$. This region follows the stellar sequence of B-type stars in the SDSS photometric system (Fukugita et al. 1996). Following Brown et al. (2006a), we impose a color cut $-0.5 < (r' - i')_0 < 0$ to reject objects with non-stellar colors. We also exclude the region of sky between $b < -l/5 + 50^\circ$ and $b > l/5 - 50^\circ$ to avoid excessive contamination from Galactic bulge stars. There are 746 SDSS DR4 candidate B stars with $15 < g'_0 < 17$ in the selection region. We have observed 514 stars, 69% of this total. The average surface number density of bright HVS candidates is 0.13 deg^{-2} . Thus we have surveyed an effective area of $\sim 4000 \text{ deg}^2$ for HVS candidates with $15 < g'_0 < 17$ with FAST.

2.2. Spectroscopic Observations and Radial Velocities

New observations at the 6.5m MMT telescope were obtained with the Blue Channel spectrograph on the nights of 2006 May 23-26 and 2006 June 19-20. The spectrograph was operated with the 832 line mm^{-1} grating in second order and a $1.25''$ slit. These settings provided a wavelength coverage of 3650 Å to 4500 Å and a spectral resolution of 1.2 Å. Exposure times ranged from 5 to 30 minutes and were chosen to yield $S/N = 15$ in the continuum at 4000 Å. Comparison lamp exposures were obtained after every exposure.

Observations at the 1.5m Tillinghast telescope were obtained with the FAST spectrograph over the course of 19 nights between 2006 January 1 and 2006 July 21. The spectrograph was operated with a 600 line mm^{-1} grating




FIG. 2.— Spectrum of the $z = 2.43$ quasar located at $15^{\text{h}}58^{\text{m}}51^{\text{s}}85, +22^{\text{d}}21^{\text{m}}59^{\text{s}}9$ (J2000).

and a $2''$ slit. These settings provided a wavelength coverage of 3500 Å to 5400 Å and a spectral resolution of 2.3 Å. Like the MMT observations, exposure times ranged from 5 to 30 minutes and were chosen to yield $S/N = 15$ in the continuum at 4000 Å. Comparison lamp exposures were obtained after every exposure.

Spectra were extracted using IRAF¹ in the standard way. Radial velocities were measured using the cross-correlation package RVSAO (Kurtz & Mink 1998). Brown et al. (2003) describe in detail the cross-correlation templates we use. The average uncertainty is $\pm 11 \text{ km s}^{-1}$ for the B-type stars.

2.3. Selection Efficiency and Completeness

Our combined observations total 894 HVS candidates with $15 < g'_0 < 19.5$, of which 699 (78%) are stars of B spectral type. The remaining objects include 185 DA white dwarfs, 3 other (DZ, DB) white dwarfs, 1 compact blue galaxy (see Kewley et al. 2006), 1 quasar, and 5 objects of uncertain identification, possibly broad absorption line quasars. The one obvious quasar is a previously unidentified $g' = 19.67$ object at $z = 2.43$, located at $15^{\text{h}}58^{\text{m}}51^{\text{s}}85, +22^{\text{d}}21^{\text{m}}59^{\text{s}}9$ (J2000). Figure 2 presents its spectrum. The quasar has colors $(u' - g')_0 = +0.80$ and $(g' - r')_0 = -0.28$, unusually red in $(u' - g')_0$ and blue in $(g' - r')_0$ compared to other quasars at similar redshifts (Schneider et al. 2005).

Figure 1 plots the spectroscopic identifications of the HVSs, other B-type stars, and the white dwarfs. Interestingly, we find stars with spectral types as early as B3 in the bright FAST sample: the solid dots in Figure 1 are the 61 stars with spectral types B3 and B4. If these early B-type stars are main sequence stars, they are located at surprisingly large heliocentric distances $30 \lesssim d \lesssim 60 \text{ kpc}$. We discuss the early B-type stars further in §3.

Our survey is now 100% complete across the high declination-region of SDSS DR4 located at $25 < l < 210^\circ$, $b > 0^\circ$ in Galactic coordinates. Curiously, we find no new unbound HVSs in the completed region. Our previous discovery rate with the MMT was 1 HVS per ~ 50 B-type stars (Brown et al. 2006b), yet our new set of observations contains 245 (57 from the MMT) B-type stars in the complete region. The absence of an unbound HVS in this large region suggests that the distribution

¹ IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

of HVSs on the sky may have structure. We address the spatial distribution of HVSs in §4.

For the remainder of this paper, we consider a “clean” sample containing 560 stars selected only from the overlapping region of color-color space: $-0.38 < (g' - r')_0 < -0.28$ and $2.67(g' - r')_0 + 1.30 < (u' - g')_0 < 2.67(g' - r')_0 + 2.0$. Main sequence B stars in this selection region have masses of 3 - 5 $M_\odot$. We also consider the full sample of 699 B-type stars at all colors. The full sample contains earlier B-type objects from the FAST sample possibly including main sequence stars with masses up to 7 $M_\odot$. We emphasize that our photometric selection is complete and can detect a B-type star at any velocity.

3. BOUND HYPERVELOCITY STARS

The observed distribution of distances and velocities of our B-type stars provides ample evidence that they are not main sequence run-aways. Curiously, the velocity distribution reveals an asymmetry of stars with large positive radial velocities. Because the large positive velocity outliers are very unlikely main sequence run-aways, we argue that the stars represent a class of *bound* HVSs ejected from the Galactic center.

3.1. Background

High Galactic latitude B-type stars were first reported by Humason & Zwicky (1947). Early spectroscopic studies (Feige 1958; Berger 1963; Greenstein 1966) showed that high latitude B-type stars are a mix of Population II post-main sequence stars and Population I main sequence run-aways. Accurate space motions and distances are now known for nearby $d < 10$ kpc B-type stars observed by *Hipparcos*. A detailed analysis by Martin (2004) shows that 2/3 of the high latitude B stars in the *Hipparcos* catalog are main sequence run-aways and 1/3 are evolved (mostly blue horizontal branch) halo stars. Martin (2004) finds that all of the main sequence run-away B stars in the *Hipparcos* catalog have orbits consistent with a disk origin.

Our HVS survey probes much fainter and more distant B-type stars than observed by *Hipparcos*. Thus we cannot rely on proper motions or parallaxes. Brown et al. (2006b) discuss the nature of the late B-type stars and conclude that the distribution of radial velocities and metallicities suggests that they are likely a Galactic halo population of post-main sequence stars and/or blue stragglers. We now re-visit the possibility of main sequence run-aways in our sample.

Theorists have shown that unbinding a stellar binary through supernova disruption (Blaauw 1961) or binary-binary interactions (Poveda et al. 1967) produces a maximum ejection velocity of 200 - 300 km s^{-1} (Leonard 1991, 1993; Portegies Zwart 2000; Davies et al. 2002; Gualandris et al. 2004; Dray et al. 2005). The maximum velocity is set by the escape velocity from the stellar surface. For example, a contact binary containing two 3 $M_\odot$ stars has a Keplerian orbital velocity of 240 km s^{-1} . Extracting a large ejection velocity from disrupting such a system is difficult. Portegies Zwart (2000) considers supernovae unbinding binaries and finds that 90% of ejections have velocities between 0 and 100 km s^{-1} ; only 1% of 3 $M_\odot$ stars are ejected at 200 km s^{-1} . Davies et al. (2002) perform similar calculations for binaries that undergo a common envelope phase followed by a mass-transfer phase




FIG. 3.— Galactic rest-frame velocity histogram of the clean sample of 560 B-type stars (*upper panel*). The best-fit Gaussian (*thin line*) has dispersion $105 \pm 5 \text{ km s}^{-1}$. The lower panel plots the residuals of the observations from the best-fit Gaussian, normalized by the value of the Gaussian. In addition to the unbound HVSs, there is a significant asymmetry of 11 positive velocity outliers $+275 < v_{rf} < 450 \text{ km s}^{-1}$ including SDSS J144955.58+310351.4.

that produces a type Ia supernova. In such systems, Davies et al. find that 3 $M_\odot$ secondaries have typical ejection velocities $\sim 100 \text{ km s}^{-1}$ and maximum ejection velocities of 250 km s^{-1} . By comparison, the escape velocity of the Galaxy near the Sun is at least 500 km s^{-1} (Carney et al. 1988).

Because main sequence run-aways travel on bound orbits, they spend most of their time near the apex of their trajectories above the disk with small line-of-sight velocities. For example, a star ejected at 240 km s^{-1} vertically out of an infinite disk with surface mass density 95 $M_\odot \text{ pc}^{-2}$ reaches an apex of $z = 11$ kpc. This simple calculation illustrates that main sequence run-aways cannot simultaneously have large velocities $v > +275 \text{ km s}^{-1}$ and large distances $z > 10$ kpc. Davies et al. (2002) use a full Galaxy potential model and compute orbits for 1000 run-away stars randomly ejected from their predicted velocity distribution. The vast majority of the Davies et al. (2002) run-aways are located at $|z| < 5$ kpc; at high Galactic latitudes $|b| > 30^\circ$ the most distant run-aways are found ~ 25 kpc from the disk.

3.2. The Asymmetric Velocity Distribution

Figure 3 plots the distribution of line-of-sight velocities, corrected to the Galactic rest-frame, for our clean sample of 560 late B-type stars plus HVS1 (Brown et al. 2005a). It is apparent from Figure 3 that the velocities are well described by a Gaussian distribution. We iteratively clip 3σ outliers and calculate a $105 \pm 5 \text{ km s}^{-1}$ dispersion and a $+14 \pm 4 \text{ km s}^{-1}$ mean for the distribution. We note that the full sample of 699 stars has a statistically identical distribution. The lower panel of Figure 3 plots the residuals of the observations from this Gaus-

sian distribution, normalized by the value of the Gaussian. Stars with velocities $|v_{rf}| < 275 \text{ km s}^{-1}$ show low-significance deviations from a Gaussian distribution and probably constitute a halo population of post-main sequence stars and/or blue stragglers (Brown et al. 2006b). The highest-significant outliers are the first HVS (HVS1), the four HVSs previously discovered in this survey (HVS4 - HVS7), and a new object: SDSS J144955.58+310351.4.

The possible HVS, SDSS J144955.58+310351.4, is a bright $g'_0 = 15.70 \pm 0.03$ star with an $A1 \pm 1.4$ spectral type and solar metallicity $[\text{Fe}/\text{H}]_{W_k} = -0.3 \pm 0.6$. A main sequence star of this type has heliocentric distance $d \simeq 17 \text{ kpc}$ (Schaller et al. 1992); an evolved blue horizontal branch (BHB) star has $d \simeq 7 \text{ kpc}$ (Clewley et al. 2005). Located at $(l, b) = (48.7^\circ, 63.9^\circ)$, the star is high above the Galactic plane at $z \simeq 15 \text{ kpc}$ and a Galactocentric distance of $R \simeq 17 \text{ kpc}$ (assuming $d = 17 \text{ kpc}$). The star's $+363 \pm 10 \text{ km s}^{-1}$ heliocentric radial velocity corresponds to a minimum velocity in the Galactic rest frame of $+447 \text{ km s}^{-1}$ (see Brown et al. 2006b). The star is listed with no proper motion in the USNOB1 (Monet et al. 2003) catalog, and thus is probably bound to the Milky Way.

Curiously, we find a set of more distant stars with equally large, but bound, velocities. Figure 3 shows a significant asymmetry in the wings of the observed velocity distribution: 11 stars (excluding the 5 unbound HVSs) have large positive velocities $+275 < v_{rf} < +450 \text{ km s}^{-1}$; only 1 star has a comparable negative velocity at $v_{rf} = -286 \pm 12 \text{ km s}^{-1}$. This asymmetry was first pointed out by Brown et al. (2006b) but is more significant in this larger sample. There is a 0.00022 probability of drawing 11 stars with $+275 < v_{rf} < 450 \text{ km s}^{-1}$ and 1 star with $-450 < v_{rf} < -275 \text{ km s}^{-1}$ from a Gaussian distributions with the observed parameters. Integrating the wings of a Gaussian with the observed parameters, we would expect to find 3.6 stars with $+275 < v_{rf} < 450 \text{ km s}^{-1}$ and 1.6 stars $-450 < v_{rf} < -275 \text{ km s}^{-1}$. Thus there is an excess of ~ 7 stars with large positive velocity and no apparent excess of stars with large negative velocity, significant at the $\sim 3.5\sigma$ level.

If the positive velocity outliers are main sequence stars, they are located at large distances 20 - 80 kpc (see Table 1). Yet the outliers cannot be run-away stars, because the run-away mechanisms cannot produce simultaneously large velocities and large distances. If, on the other hand, the positive velocity outliers are halo stars on radial orbits, we would expect to find equal numbers of stars moving toward and away from us, contrary to observation. Compact binary systems may also produce outliers in the velocity distribution. But velocity outliers resulting from compact binaries should be distributed symmetrically, again contrary to observation. We note that neutron stars are observed traveling at velocities of 1000 km s^{-1} , but such neutron stars are the remnants of asymmetric supernova explosions, and are not B-type stars.

3.3. Bound HVSs

One explanation for the significant excess of B-type stars traveling $+275 < v_{rf} < 450 \text{ km s}^{-1}$ is the HVS mechanism. HVSs are ejected by three-body interactions involving a MBH (Hills 1988). HVSs almost cer-




FIG. 4.— Predicted velocity distributions of 3-5 $M_\odot$ HVSs with distance from the Galactic center (Bromley et al. 2006). We calculate distances for our B-type stars assuming they are main sequence stars: the four HVSs discovered by this survey (solid squares) and the 11 positive velocity outliers $+275 < v_{rf} < 450 \text{ km s}^{-1}$ (open squares) including SDSS J144955.58+310351.4 (open square with plus sign).

tainly come from the Galactic Center. The presence of a $3.6 \times 10^6 M_\odot$ MBH (Ghez et al. 2005; Eisenhauer et al. 2005) in the crowded Galactic center inevitably produces a “fountain” of HVSs ejections. The actual HVS ejection rate depends on the number of stars with radial orbits in the MBH’s so-called “loss cone.” If the MBH’s loss cone is replenished with stars scattered in by massive star clusters, molecular clouds, or IMBHs (Perets et al. 2006), then the ejection rate can be orders of magnitude larger than the original Yu & Tremaine (2003) predictions. Regardless of the exact rate, this fountain picture results in a broad spectrum of HVS ejection velocities which include bound and unbound orbits (Hills 1991; Yu & Tremaine 2003; Gualandris et al. 2005; Levin 2006; Baumgardt et al. 2006; Perets et al. 2006; Sesana et al. 2006; Bromley et al. 2006; O’Leary & Loeb 2006). Bromley et al. (2006) recently calculated ejection velocities for binaries containing 3-4 $M_\odot$ primaries matched to our survey of B-type stars.

Over the volume sampled by our survey, Bromley et al. (2006) predict comparable numbers of HVSs ejected into bound and unbound orbits. Our survey has discovered 4 HVSs on unbound orbits suggesting that 4 ± 2 of our excess ~ 7 positive velocity outliers are plausibly HVSs on bound orbits. Although the predicted and observed numbers of bound HVSs are statistically consistent, we note that additional HVS mechanisms may be at work. For example, O’Leary & Loeb (2006) show that single stars can be ejected by encounters with stellar mass black holes orbiting the central MBH. Such HVSs tend to have lower ejection velocities than HVSs ejected by the Hills (1988) mechanism, and thus may account for additional HVSs on bound orbits.

Figure 4 plots the Bromley et al. (2006) predicted velocity distributions for main sequence HVSs ejected by the central MBH as a function of Galactocentric radius. HVS velocities increase with depth because only the most rapidly moving HVSs survive long enough to reach large Galactocentric distances. The large squares in Figure 4 are our 4 HVSs (solid squares) and the 11 stars with $+275 < v_{rf} < 450 \text{ km s}^{-1}$ (open squares). We assume that the stars are main-sequence stars, and use the observed colors and spectral types of the stars to estimate

stellar luminosity, distance, and stellar mass (see Table 1). The solid lines in Figure 4 indicate our survey limits for main sequence stars of a given stellar mass.

We evaluate the probability of drawing our observed sample from the Bromley et al. (2006) predicted distributions. We use the two-dimensional, two-sample Kolmogorov-Smirnov test (Press et al. 1992) to compare the observed and predicted distributions of HVS velocity and distance. First, we generate 10^5 realizations of the observations – 4 HVSs plus 7 bound HVSs randomly drawn from the 11 high velocity outliers – and calculate the maximum difference in integrated probability against the Bromley et al. (2006) distributions. We use only the portion of the Bromley et al. (2006) distributions that fall within our observational survey limits (see Figure 4). Second, we randomly sample 11 objects from the Bromley et al. (2006) distributions 10^5 times and calculate the maximum difference in integrated probabilities against the Bromley et al. (2006) distributions. Again, we use only the portion of the Bromley et al. (2006) distributions that fall within our observational survey limits. Finally, the likelihood is the fraction of time that the probabilities of the synthetic data sets exceed the probabilities of the real data sets. The mean likelihood of drawing our observed sets of stars from the predicted distributions is approximately 0.128, which neither strongly supports nor strongly rejects the Bromley et al. (2006) model. We conclude that the most plausible explanation for the observed excess of positive velocity outliers is the HVS mechanism ejecting stars from the Galactic center on bound orbits.

We note that other stars in the literature may also be explained as bound HVSs. For example, the $5 M_{\odot}$ main sequence B star HIP 60350 has a heliocentric velocity of $+230 \text{ km s}^{-1}$ and a full space velocity of 417 km s^{-1} based on *Hipparcos* measurements (Maitzen et al. 1998). The star’s large radial motion $U = -352 \text{ km s}^{-1}$ means that it originated from well inside the solar circle, perhaps from the Galactic center.

3.4. Mystery of the Early B-type Stars

Our full sample of B-type stars contains many early-type stars at bright magnitudes. Four of the early B stars in the full sample are possibly bound HVSs with $+275 < v_{rf} < 400 \text{ km s}^{-1}$. If they are main sequence stars, the four early B stars have masses of $5 - 7 M_{\odot}$ and they are located at distances ranging 30 - 60 kpc (bounded by the FAST survey limiting magnitudes). We now ask whether the number of early and late B-type bound HVSs are consistent with a standard initial mass function. Assuming the stars are main sequence stars, there are nine $3 - 5 M_{\odot}$ stars in our sample located in the same volume as the four $5 - 7 M_{\odot}$ stars: a ratio of 2.25. A Salpeter initial mass function predicts a ratio of 2.7, similar to what we observe. However, the lifetimes of $5 - 7 M_{\odot}$ main sequence stars are only 40 - 90 Myr (Schaller et al. 1992), a factor of 2 - 3 shorter than $3 - 5 M_{\odot}$ stars. The travel times of the four early B stars, assuming v_{rf} is their full space velocity, range from ~ 100 to ~ 150 Myr from the Galactic center. Thus if the early B stars are main sequence stars ejected from the Galactic center, they do not have high enough velocities to survive to their inferred distances.

How can we reconcile the lifetimes of the early B-type

TABLE 1
POSSIBLE BOUND HVSs

Catalog ID	l deg	b deg	v_{rf} km s^{-1}	type	R^a kpc	R^b kpc
SDSS J074950.24+243841.2	196.1	23.2	+293	B8	66	27
SDSS J075055.24+472822.9	171.5	29.4	+307	B6	42	17
SDSS J075712.93+512938.0	167.0	30.9	+329	B7	32	13
SDSS J081828.07+570922.1	160.4	34.2	+283	B9	42	24
SDSS J090710.08+365957.5	186.3	42.2	+280	B9	57	29
SDSS J110224.37+025002.8	251.2	54.4	+326	A1	51	23
SDSS J115245.91-021116.2	274.9	57.5	+305	A1	53	18
SDSS J140432.38+352258.4	65.3	72.4	+293	B8	51	18
SDSS J141723.34+101245.7	357.2	63.6	+289	A1	48	19
SDSS J142001.94+124404.8	2.5	64.8	+300	B9	27	12
SDSS J144955.58+310351.4	48.7	63.9	+447	A1	16	9

^aGalactocentric distance for a main sequence star of the observed spectral type.

^bGalactocentric distance for a BHB star of the observed color.

stars with their travel times from the Galactic center? It is possible that the early B-type stars are blue stragglers, main sequence stars that have undergone mass transfer or mergers. Blue stragglers are usually associated with globular clusters, yet surveys of field BHB stars in the halo find that half of the stars are in fact high surface-gravity blue stragglers or A dwarfs (Norris & Hawkins 1991; Preston et al. 1994; Wilhelm et al. 1999; Brown et al. 2003; Clewley et al. 2004; Brown et al. 2005b). Because HVSs likely originate in tight binary systems, it is possible that some HVSs experience mass-transfer from their former companion prior to being ejected from the Galactic center. A blue straggler origin is one explanation for HVS2, an $8 M_{\odot}$ HVS with a main sequence lifetime otherwise inconsistent with travel time from the Galactic center (Edelmann et al. 2005).

It is also possible that the early B-type stars are main sequence stars ejected elsewhere in the Galaxy by dynamical interactions with compact objects like intermediate mass black holes (IMBHs). However, as the mass of a black hole decreases, the cross-section of HVS interaction decreases (e.g. Hills 1988). A stellar binary must come much closer to an IMBH before the gravitational tidal energy exceeds the binding energy of the binary. Thus an IMBH produces fewer HVSs than a MBH in the same environment. Unless IMBHs are ubiquitous in dense stellar environments such as the centers of globular clusters, we expect that the overall HVS rate is dominated by the MBH in the Galactic center.

A final possibility is that the early B-type stars are post main-sequence stars. Demarque & Virani (2007) argue that the Galactic center contains many more old dwarf stars than young B stars, and thus the MBH should eject many more BHB stars than main sequence B stars. Davies & King (2005) and Dray et al. (2006) argue that the Galactic center contains tidally-stripped evolved stars (in effect, BHB stars with small hydrogen envelopes) that mimic B stars. B-type BHB stars with hot effective temperatures and small hydrogen envelopes spend most of their helium-burning lifetimes at intrinsically faint luminosities (Yi et al. 1997; Dray et al. 2006). Thus, if they are BHB stars, the early B-type stars are located 10 - 20 kpc from the Galactic center and have




FIG. 5.— Aitoff sky map, in Galactic coordinates, showing the clean sample of 560 B-type HVS candidates. Radial velocities, in the Galactic rest frame, are indicated by the color of the solid squares: purple is -300 , green is 0 , and red is $+300$ km s^{-1} . The HVSs are completely off this color scale and are marked by: HVS1 (plus), the four HVSs discovered by this survey (solid squares), and the 11 positive velocity outliers $+275 < v_{r,f} < 400$ km s^{-1} (open squares) including SDSS J144955.58+310351.4 (open square with plus sign). Dotted lines are lines of constant unbound HVS fraction calculated from the Bromley et al. (2006) models.

travel times ranging from 25 - 50 Myr. The BHB progenitor lifetimes easily exceed the inferred travel times, reconciling any difficulties with the MBH ejection origin. To test the post-main sequence explanation requires high dispersion spectroscopy to determine the stellar nature of the bound HVSs.

4. HVS SPATIAL ANISOTROPY

The distribution of HVSs on the sky is interesting because the spatial distribution of HVSs is linked to their origin. Bromley et al. (2006) show that fraction of unbound HVSs increases with radial distance from the Galactic center (see Figure 4). The fraction changes with depth because only the most rapidly moving HVSs survive to reach large Galactocentric distances. Because the Sun is located 8 kpc from the Galactic center, our survey reaches approximately 16 kpc deeper towards the anti-center than towards the Galactic center. Thus, if the fountain model of HVSs ejected from the Galactic center is correct, we expect to find more unbound HVSs towards the anti-center.

We use the Bromley et al. (2006) models (Figure 4) to calculate the expected fraction of unbound HVSs in our survey volume as a function of position on the sky. The predicted fraction of unbound HVSs ranges from 53% towards the Galactic center to 63% towards the anti-center. The fraction does not vary much because it is dominated by the survey volume at large distances. The HVSs found in this survey, however, are not located at distances beyond ~ 80 kpc. If we limit the survey volume to the observed range of HVS distances, then the predicted fraction of unbound HVSs ranges from 31% to 46%. We now consider the observations.

Figure 5 plots the distribution of our clean sample of 560 B-type stars across the sky in Galactic coordinates. Color indicates the radial velocity, in the Galac-

tic rest frame, of the stars. The four unbound HVSs (solid black squares) discovered by this survey are completely off the color scale; HVS1 is marked by a plus sign. The 11 possible bound HVSs are indicated by open squares; SDSS J144955.58+310351.4 is plotted with an open square with a plus sign. The dotted lines in Figure 5 are lines of constant HVS fraction calculated from the Bromley et al. (2006) models. We note that our survey is currently incomplete in the SDSS equatorial slices, located at $l > 210^\circ$ and $b < 0^\circ$.

It is striking that all of the unbound HVSs in our survey are located towards the anti-center. We randomize the observed velocities among the positions of all 560 stars, and find a 0.064 probability of randomly drawing the 4 unbound HVSs from our survey in the hemisphere $90 < l < 270^\circ$. Thus the unbound HVSs have a marginally anisotropic spatial distribution, being preferentially located in the anti-center hemisphere with $\sim 2\sigma$ confidence. We note that the 3 unbound HVSs not from this survey, HVS1 (Brown et al. 2005a), HVS2 (Edelmann et al. 2005), and HVS3 (Hirsch et al. 2005), are also located in the anti-center hemisphere $90 < l < 270^\circ$.

The 11 possible bound HVSs, on the other hand, appear more evenly distributed across the sky: 6 bound HVSs are located $90 < l < 270^\circ$ and 5 bound HVSs are located $|l| < 90^\circ$. To calculate the variation in unbound HVS fraction, we consider only 7 bound HVSs (the excess number of high velocity outliers in Figure 3) and split them 4/3 between the anti-center and Galactic-center hemispheres, respectively. The fraction of unbound HVSs is then $50 \pm 30\%$ in the anti-center hemisphere and $0 \pm 40\%$ in the Galactic-center hemisphere. Although small number statistics overwhelm our measurement, the unbound HVS fraction appears consistent with the predictions of Bromley et al. (2006) and the Galactic center ejection picture.

There is another possible explanation for spatial anisotropy: HVSs ejected via single star encounters with a MBH binary are preferentially ejected in the orbital plane of the MBH binary (Gualandris et al. 2005; Holley-Bockelmann et al. 2005; Levin 2006; Sesana et al. 2006; Merritt 2006; Baumgardt et al. 2006). Recent N-body simulations for the case of an IMBH inspiralling into a MBH, however, show that HVSs are ejected nearly isotropically because the angular momentum vector of the IMBH rapidly changes on a ~ 1 Myr timescale (Baumgardt et al. 2006). Moreover, our observed set of HVSs do not share common travel times from the Galactic center. Different IMBH inspiral events would likely have different orbital histories, thus it seems unlikely that the observed HVS spatial distribution can point back to a binary MBH origin. A set of HVSs with common travel times must be found in order to test the binary MBH origin.

5. CONCLUSIONS

We discuss our targeted survey for HVSs, a spectroscopic survey of stars with B-type colors. We introduce a new component of the survey that samples brighter and bluer stars using the FAST spectrograph. Our combined set of observations now contains 699 stars of B spectral type, 188 white dwarfs, and 1 previously unidentified $z = 2.43$ quasar located at $15^{\text{h}}58^{\text{m}}51^{\text{s}}.85, +22^{\circ}21'59''.9$ (J2000).

We find no new unbound HVS, but we find evidence for a new class of *bound* HVSs. The velocity distribution of the B-type stars shows a significant excess of ~ 7 stars with large positive velocities $+275 < v_{rf} < 450 \text{ km s}^{-1}$ and no apparent excess of stars with large negative velocity, significant at the $\sim 3.5\sigma$ level. Neither halo stars on radial orbits, compact binary stars, nor main sequence run-away stars ejected from the disk can explain simultaneously the large velocities and distances of these stars. The most plausible explanation is that the positive velocity outliers are HVSs ejected from the Galactic center on bound orbits.

Our bright FAST survey contains a large number of early B-type stars, four of which have velocities $+275 < v_{rf} < 450 \text{ km s}^{-1}$. These four possible bound HVSs have main sequence lifetimes too short to survive to their inferred distances, however. One explanation is that the stars are post-main sequence stars. Establishing HVSs as main sequence or post-main sequence stars is important for measuring the stellar mass function of HVSs and probing the types of stars that orbit near the MBH.

Stellar rotation is a useful discriminant between rapidly rotating main sequence B stars (Abt et al. 2002; Martin 2004) and slowly rotating BHB stars (Peterson et al. 1995; Behr 2003). Metallicity is also a good discriminant between young main sequence B stars and post-main sequence halo stars. Echelle observations of the brightest HVSs, underway now, will reveal the stars' true nature.

A Galactic center origin predicts that unbound HVSs should be found preferentially towards the anti-center as a result of survey selection. We observe that the unbound HVSs have a marginally anisotropic distribution at $2\text{-}\sigma$ confidence. Our estimates of unbound HVS fraction are consistent with the Galactic center ejection picture. Measuring the spatial distribution of a larger sample of HVSs may provide a strong test of the stars' origin.

We note that full space motions for the HVSs will be known in a few years. A *Hubble Space Telescope* program is measuring the stars' positions with 0.5 mas accuracies by registering background galaxies across Advanced Camera for Surveys images. The HVSs have predicted proper motions of $0.5 - 1 \text{ mas yr}^{-1}$, thus 3σ proper motion measurements are possible over a three year baseline (Gnedin et al. 2005).

We are continuing our targeted HVS survey of faint B-type stars. SDSS Data Release 5 provides additional targets that we are now following-up with the MMT and Whipple 1.5m telescopes. Given our current discovery rate, we expect to find another few unbound HVSs in the coming months. Identifying HVSs throughout the Galaxy will allow us to measure the mass function of stars in the Galactic center and the history of stellar interactions with the central MBH.

We thank M. Alegria, J. McAfee, and A. Milone for their assistance with observations obtained at the MMT Observatory, a joint facility of the Smithsonian Institution and the University of Arizona. We thank H. Perets for helpful correspondence. This project makes use of data products from the Sloan Digital Sky Survey, which is managed by the Astrophysical Research Consortium for the Participating Institutions. This research has made use of NASA's Astrophysics Data System Bibliographic Services. This work was supported by W. Brown's Clay Fellowship and the Smithsonian Institution.

Facilities: MMT (Blue Channel Spectrograph), FLWO:1.5m (FAST Spectrograph)

REFERENCES

- Abt, H. A., Levato, H., & Grosso, M. 2002, *ApJ*, 573, 359
 Adelman-McCarthy, J. K. et al. 2006, *ApJS*, 162, 38
 Baumgardt, H., Gualandris, A., & Portegies Zwart, S. 2006, *MNRAS*, 372, 174
 Behr, B. B. 2003, *ApJS*, 149, 67
 Berger, J. 1963, *PASP*, 75, 393
 Blaauw, A. 1961, *Bull. Astron. Inst. Netherlands*, 15, 265
 Bromley, B. C., Kenyon, S. J., Geller, M. J., Barcikowski, E., Brown, W. R., & Kurtz, M. J. 2006, *ApJ*, 653, 1194
 Brown, W. R., Allende Prieto, C., Beers, T. C., Wilhelm, R., Geller, M. J., Kenyon, S. J., & Kurtz, M. J. 2003, *AJ*, 126, 1362
 Brown, W. R., Geller, M. J., Kenyon, S. J., & Kurtz, M. J. 2005a, *ApJ*, 622, L33
 —. 2006a, *ApJ*, 640, L35
 —. 2006b, *ApJ*, 647, 303
 Brown, W. R., Geller, M. J., Kenyon, S. J., Kurtz, M. J., Allende Prieto, C., Beers, T. C., & Wilhelm, R. 2005b, *AJ*, 130, 1097
 Carney, B. W., Laird, J. B., & Latham, D. W. 1988, *AJ*, 96, 560
 Clewley, L., Warren, S. J., Hewett, P. C., Norris, J. E., & Evans, N. W. 2004, *MNRAS*, 352, 285
 Clewley, L., Warren, S. J., Hewett, P. C., Norris, J. E., Wilkinson, M. I., & Evans, N. W. 2005, *MNRAS*, 362, 349
 Davies, M. B. & King, A. 2005, *ApJ*, 624, L25
 Davies, M. B., King, A., & Ritter, H. 2002, *MNRAS*, 333, 463
 Demarque, P. & Virani, S. 2007, *A&A*, 461, 651
 Dray, L. M., Dale, J. E., Beer, M. E., Napiwotzki, R., & King, A. R. 2005, *MNRAS*, 364, 59
 Dray, L. M., King, A. R., & Davies, M. B. 2006, *MNRAS*, 910
 Edelmam, H., Napiwotzki, R., Heber, U., Christlieb, N., & Reimers, D. 2005, *ApJ*, 634, L181

- Eisenhauer, F. et al. 2005, *ApJ*, 628, 246
 Fabricant, D., Cheimets, P., Caldwell, N., & Geary, J. 1998, *PASP*, 110, 79
 Feige, J. 1958, *ApJ*, 128, 267
 Fuentes, C. I., Stanek, K. Z., Gaudi, B. S., McLeod, B. A., Bogdanov, S., Hartman, J. D., Hickox, R. C., & Holman, M. J. 2006, *ApJ*, 636, L37
 Fukugita, M., Ichikawa, T., Gunn, J. E., Doi, M., Shimasaku, K., & Schneider, D. P. 1996, *AJ*, 111, 1748
 Ghez, A. M., Salim, S., Hornstein, S. D., Tanner, A., Lu, J. R., Morris, M., Becklin, E. E., & Duchene, G. 2005, *ApJ*, 620, 744
 Ginsburg, I. & Loeb, A. 2006a, *MNRAS*, 368, 221
 —. 2006b, preprint astro-ph/0609440
 Gnedin, O. Y., Gould, A., Miralda-Escudé, J., & Zentner, A. R. 2005, *ApJ*, 634, 344
 Greenstein, J. L. 1966, *ApJ*, 144, 496
 Gualandris, A. & Portegies Zwart, S. 2007, *MNRAS*, accepted
 Gualandris, A., Portegies Zwart, S., & Eggleton, P. P. 2004, *MNRAS*, 350, 615
 Gualandris, A., Portegies Zwart, S. P., & Sipior, M. S. 2005, *MNRAS*, 363, 223
 Hills, J. G. 1988, *Nature*, 331, 687
 —. 1991, *AJ*, 102, 704
 Hirsch, H. A., Heber, U., O'Toole, S. J., & Bresolin, F. 2005, *A&A*, 444, L61
 Holley-Bockelmann, K., Sigurdsson, S., Mihos, C. J., Feldmeier, J. J., Ciardullo, R., & McBride, C. 2005, preprint astro-ph/0512344
 Humason, M. L. & Zwicky, F. 1947, *Contributions from the Mount Wilson Observatory / Carnegie Institution of Washington*, 724, 1
 Kewley, L. J., Brown, W. R., Geller, M. J., Kenyon, S. J., & Kurtz, M. J. 2006, *AJ*, accepted
 Kollmeier, J. A. & Gould, A. 2007, preprint astro-ph/0701350
 Kurtz, M. J. & Mink, D. J. 1998, *PASP*, 110, 934
 Leonard, P. J. T. 1991, *AJ*, 101, 562
 Leonard, P. J. T. 1993, in *ASP Conf. Ser.* 45, *Luminous High-Latitude Stars*, ed. D. D. Sasselov, 360
 Levin, Y. 2006, *ApJ*, 653, 1203
 Maitzen, H. M., Paunzen, E., Pressberger, R., Slettebak, A., & Wagner, R. M. 1998, *A&A*, 339, 782
 Martin, J. C. 2004, *AJ*, 128, 2474
 Merritt, D. 2006, *ApJ*, 648, 976
 Monet, D. G. et al. 2003, *AJ*, 125, 984
 Norris, J. E. & Hawkins, M. R. S. 1991, *ApJ*, 380, 104
 O'Connell, W. O. R. 1973, *AJ*, 78, 1074
 O'Leary, R. M. & Loeb, A. 2006, preprint astro-ph/0609046
 Perets, H. B., Hopman, C., & Alexander, T. 2006, preprint astro-ph/0606443
 Peterson, R. C., Rood, R. T., & Crocker, D. A. 1995, *ApJ*, 453, 214
 Portegies Zwart, S. F. 2000, *ApJ*, 544, 437
 Poveda, A., Ruiz, J., & Allen, C. 1967, *Bol. Obs. Tonantzintla Tacubaya*, 4, 860
 Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P. 1992, *Numerical recipes in C. The art of scientific computing* (Cambridge: University Press, 2nd ed.)
 Preston, G. W., Beers, T. C., & Shectman, S. A. 1994, *AJ*, 108, 538
 Schaller, G., Schaerer, D., Meynet, G., & Maeder, A. 1992, *A&AS*, 96, 269
 Schneider, D. P. et al. 2005, *AJ*, 130, 367
 Sesana, A., Haardt, F., & Madau, P. 2006, *ApJ*, in press
 Wilhelm, R., Beers, T. C., Sommer-Larsen, J., Pier, J. R., Layden, A. C., Flynn, C., Rossi, S., & Christensen, P. R. 1999, *AJ*, 117, 2329
 Worthey, G., Faber, S. M., Gonzalez, J. J., & Burstein, D. 1994, *ApJS*, 94, 687
 Yi, S., Demarque, P., & Kim, Y.-C. 1997, *ApJ*, 482, 677
 Yu, Q. & Tremaine, S. 2003, *ApJ*, 599, 1129

APPENDIX

DATA TABLE

Table 2 [on-line data table, attached as tab2.dat in the source] lists the 643 new observations from our HVS survey, excluding the extra-galactic objects. 134 HVS candidates were observed with the MMT and 509 were observed with FAST (see §2). Table 2 includes columns of RA and Dec coordinates (J2000), g' apparent magnitude, $(u' - g')_0$ and $(g' - r')_0$ color, and our heliocentric velocity v_{helio} . The column WD indicates whether the object is a B-type star (WD=0) or a white dwarf (WD=1).

We note that $(u' - g')_0$ color correlates strongly with the observed spectral type of our B-type stars. Our spectral types are completely independent of the SDSS photometry. We classify spectral types based on O'Connell (1973) and Worthey et al. (1994) line indices as described in Brown et al. (2003). Typical uncertainty is ± 1.6 spectral sub-types. Figure A6 plots our spectral types and the $(u' - g')_0$ color for our full sample of 699 B-type stars. The solid line is the best-fit relation: spectral type = $9.5(u' - g')_0 + 11.5$, where spectral type 10=B0, 15=B5, 20=A0, etc. The spectral types of the stars in our HVS survey range B3 - A2, with a dispersion around the best-fit relation of ± 1 spectral sub-types or ± 0.1 mag in $(u' - g')_0$.




FIG. A6.— Spectral types determined from our spectroscopy and SDSS $(u' - g')_0$ color for the full sample of 699 B-type stars.